

Audit and feedback + educational meetings versus no intervention - dichotomous outcomes															
Study	Outcome	N health professional	Ctr before	A&F before	Ctr after	A&F after	RD Unadjust	RDAdjust	RR Unadjust	RR Adjust	Baseline compliance	Study quality	Intensity of A&F	Complexity of targeted behaviour	Seriousness of outcome
De Almeida Neto 2000	% analgesic misuse identified and discussed (mean of two outcomes)	22					27.00	24.00	4.41	3.01	6%	moderate	moderate	moderate	moderate
Veninga 1999a	% correct prescribing for asthma -Netherlands (median of four outcomes)	181					4.50	5.50	1.10	1.09	65%	moderate	moderate	moderate	moderate
Veninga 1999b	% correct prescribing fro asthma -Sweden (median of four outcomes)	204					2.50	-1.00	1.04	0.98	51%	moderate	moderate	moderate	moderate
Veninga 1999c	% compliance with guidelines for asthma -Norway (median of four outcomes)	199					2.50	1.50	1.05	1.03	63%	moderate	moderate	moderate	moderate
Veninga 1999d	% compliance with guidelines for asthma -Slovakia (median of four outcomes)	81					-0.50	1.00	1.00	1.06	40%	moderate	moderate	moderate	moderate