

Multifaceted interventions that include audit and feedback versus no intervention	Audit and feedback as part of multifaceted intervention no intervention - continuous outcomes													
Study	Outcome	N health professionals	Ctr before	A&F before	Ctr after	A&F after	Post difference in desired direction	Adjust change in desired direction	Prosent change relative to control after	Baseline value	Study quality	Intensity of A & F	Comp-lexity of targeted behaviour	Serious-ness of targeted behaviour
van der Weijden 1999	OR for appropriate cholesterol case finding	32	4.2	5.1	3.8	4.9	1.10	0.2	5%	5.1	moderate	moderate	moderate	moderate
McConnell 1982	Median prescription of tetracycline for upper respiratory tract infaction (median of sixteen outcomes)	33					2.00	1.25	60%	7.25	moderate	moderate	low	moderate
Zwar IBD	Antibiotic prescriptions for upper respiratory infections per 100 diagnosis	157	22	25	31.7	19.7	12.00	15	47%	25	moderate	moderate	moderate	moderate
Eccles	Requests per 1000 of knee and lumbar spine radiographs (mean of two outcomes)	40					1.69	3.91	49%	8.82	moderate	low	low	low
Finkelstein 2001	Total number of antimicrobials dispensed divided by total number of person-year (mean of two outcomes)	157					0.6	0.06	3%	1.82	moderate	moderate	low	low
Verstappen 2003 IBD a	Mean number of inappropriate tests, per physician per 6 months	174	66	63	63	45	18.00	15	24%	63	high	moderate	low	low
Verstappen 2003 IBD b	Mean number of inappropriate tests, per physician per 6 months	174	134	163	126	138	-12.00	17	13%	163	high	moderate	low	low